arXiv:1210.2408v1 [cond-mat.supr-con] 8 Oct 2012

Superconductivity at the onset of spin-density-wave order in a metal

Yuxuan Wang and Andrey V. Chubukov[1]

[1] *Department of Physics, University of Wisconsin-Madison, Madison, WI 53706, USA*

Abstract

We revisit the issue of superconductivity at the quantum-critical point between a 2D paramagnet and a spin-density-wave (SDW) metal with ordering momentum (π, π). This problem is highly non-trivial because the system at criticality displays a non-Fermi liquid behavior and because the effective coupling constant λ for the pairing is generally of order one, even when the actual interaction is smaller than fermionic bandwidth. Previous study [M. A. Metlitski, S. Sachdev, Phys.Rev.B 82, 075128 (2010)] has found that the leading renormalization of the pairing vertex contains $\log^2$, like in color superconductivity. We analyze the full gap equation and argue that summing up $\log^2$ term does not lead to a pairing instability. Yet, superconductivity has no threshold, even if λ is set to be small: the subleading log terms give rise to BCS-like $T_c \propto e^{-1/\lambda}$. We argue that the analogy with BCS is not accidental as superconductivity at a QCP is a Fermi liquid phenomenon – it comes from fermions which retain Fermi liquid behavior at criticality. We computed T_c for the actual λ and found a good agreement with the numerical results.

Introduction. Superconductivity at the onset of density-wave order in a metal is an issue of high current interest, with examples ranging from cuprates [1], to Fe-pnictides [2] and other correlated materials [3–5] It is widely believed that the pairing in these systems is caused by repulsive electron-electron interaction, enhanced in a particular spin or charge channel, which becomes critical at the quantum-critical point (QCP). The pairing problem at QCP is highly non-trivial in $D \leq 3$, as scattering by a critical collective mode destroys Fermi liquid (FL) behavior above T_c (Ref. [6,9]). This is particularly relevant for systems near uniform density-wave instability (e.g., a ferromagnetic or a nematic one). In this case, FL behavior is lost on the whole Fermi-surface (FS), and superconductivity can be viewed as a pairing of incoherent fermions which exchange quanta of gapless collective bosons [7–12]. The pairing of incoherent fermions is qualitatively different from BCS/Eliashberg pairing of coherent fermions in a FL because in the incoherent case the pairing in $D < 3$ occurs only if the interaction exceeds a certain threshold [8,13,14]. For $D = 3$ there is no threshold, but at small coupling constant λ, $\log \Lambda/T_c = 1/\sqrt{\lambda}$ rather than $1/\lambda$ (Ref. [16]), in close analogy to T_c in color superconductivity (CSC) of quarks mediated by the exchange of gluons [17]

The non-FL behavior at criticality is less pronounced for systems near density-wave order at a finite momentum, because only fermions near particular points along the FS (hot spots) lose FL behavior at criticality. Still, fermions from hot regions mostly contribute to the pairing, and early studies of superconductivity at the onset of (π, π) spin-density-wave (SDW) order [8,9] placed the pairing problem into the same universality class as for QCP with $q = 0$. The 2D problem has been recently re-analyzed [19] by Metlitski and Sachdev (MS). They argued that it is important to include into the consideration the momentum dependence of the self-energy along the FS, neglected in earlier studies. Using the full form of $\Sigma(\omega_m, \mathbf{k})$ for $\mathbf{k}$ on the FS, they found that the one-loop renormalization of the pairing vertex is larger than previously thought – it is $\log^2$ instead of log, and that the enhancement comes from fermions somewhat away from hot spots, for which $\Sigma(\omega_m, \mathbf{k})$ has a FL form at the smallest frequencies. The $\log^2$ behavior in the perturbation theory holds for CSC, and MS result raises the question whether the pairing problem at a 2D SDW QCP is in the same universality class as CSC. The related issues raised by MS work are: (i) is the problem analogous to the pairing at a 2D SDW QCP a FL phenomenon, or non-FL physics is essential, (ii) what sets the scale of T_c, and (iii) is T_c non-zero only if the coupling λ exceed a finite threshold, as it happens if one approximates $\Sigma(\omega_m, \mathbf{k})$ by $\Sigma(\omega_m)$ at a hot spot, or T_c

is non-zero even at smallest λ, like in CSC?

In this letter, we address these issues. We first show that the analogy with CSC does not extend beyond one-loop order, and in our case the summation of $\log^2$ terms in the Cooper channel does not give rise to a pairing instability. However, that subleading log terms do give rise to a pairing instability, and at weak coupling yield $\log \Lambda/T_c = 1/\lambda$, like in BCS theory. We show that the analogy with BCS formula is not accidental because the pairing predominantly comes from fermions for which fermionic self-energy has a FL form. In this respect, non-FL physics at a QCP is actually irrelevant to the pairing. We then analyze the physical case $\lambda = O(1)$ in the large N approximation, assuming that terms with $n \geq 6$ loops, which break $1/N$ expansion [18,19,21], do not change the analysis qualitatively. We argue that the scale of T_c is set by the frequency ω_0 at which $\Sigma(\omega_m)$ at a hot spot becomes equal to ω_m. The theory is under control when $\omega_0 < v_F/a$ (a is inter-atomic spacing), which in practice implies that the interaction must be smaller than the fermionic bandwidth. We show that the prefactor c in $T_c = c\omega_0$ depends on the geometry of the FS and is $c = 0.04$ when the velocities at the two hot spots separated by (π, π) are nearly orthogonal. This number agrees with the slope of T_c obtained by solving the gap equation numerically along the full FS [22].

The model. We follow earlier works [8,9,19] and consider a system of fermions with N flavors and a large FS, like in the cuprates. Fermions are interacting by exchanging quanta of collective excitations in the spin channel. The static propagator of collective spin excitations is created by fermions with energies of order bandwidth and is an input for the low-energy theory of superconductivity. We assume that collective spin excitations are peaked at $\mathbf{Q} = (\pi, \pi)$ and focus on the hot regions on the FS, for which $\mathbf{k}_F + \mathbf{Q}$ is also near the FS. The Lagrangian of the model is given by [3,9,19]

$$\begin{aligned} \mathcal{S} \;=\; & -\int_k^\Lambda G_0^{-1}\left(k\right)\psi^\dagger_{k,\alpha}\psi_{k,\alpha} + \frac{1}{2}\int_q^\Lambda \chi_0^{-1}\left(q\right)\;\mathbf{S}_q\cdot\mathbf{S}_{-q} \\ & + g\int_{k,q}^\Lambda \psi^\dagger_{k+q,\alpha}\sigma_{\alpha\beta}\psi_{k,\beta}\cdot\mathbf{S}_{-q}. \end{aligned} \tag{1}$$

where $\int_k^\Lambda$ stands for the integral over $d-$dimensional $\mathbf{k}$ (up to some upper cutoff Λ) and the sum over fermionic and bosonic Matsubara frequencies, $G_0\left(k\right) = G_0(\omega_m, \mathbf{k}) = 1/(i\omega_m - \mathbf{v}_{F,\mathbf{k}}(\mathbf{k} - \mathbf{k}_F))$ is the bare fermion propagator, and $\chi_0\left(q\right) = \chi_0(\Omega_m, \mathbf{q}) = \chi_0/(\mathbf{q}^2 + \xi^{-2})$ is the static propagator of collective bosons, in which ξ^{-1} measures a distance to a QCP and $\mathbf{q}$ is measured with respect to $\mathbf{Q}$. We set $\xi^{-1} = 0$ below. The fermion-boson coupling g and χ_0 appear in theory only in combination $\bar{g} = g^2\chi_0$ and we will use $\bar{g}$ below. The Fermi velocities at hot spots separated by $\mathbf{Q}$ can be expressed as $\mathbf{v}_{F,1} = (v_x, v_y)$ and $\mathbf{v}_{F,2} = (-v_x, v_y)$, where x axis is along $\mathbf{Q}$. We will also use $\alpha = v_y/v_x$ and $v_F = (v_x^2 + v_y^2)^{1/2}$. The model of Eq. (1) can be equivalently viewed as a four-patch model for fermions near hot spots at $\pm\mathbf{k}_F$ and $\pm(\mathbf{k}_F + \mathbf{Q})$ (Ref. [19,21]). The hot spot model is obviously justified only when the interaction $\bar{g}$ is smaller than E_F.

The fermion-boson coupling gives rise to fermionic and bosonic self-energies. In the normal state, bosonic self-energy accounts for Landau damping of spin excitations, while fermionic self-energy accounts for the mass renormalization and a finite lifetime of a fermion. At one-loop level, self-consistent normal-state analysis yields [9,19,20]

$$\chi(\Omega_m, \mathbf{q}) = \frac{\chi_0}{\mathbf{q}^2 + |\Omega_m|\gamma} \tag{2}$$

$$\Sigma(\omega_m, \mathbf{k}_F) = \frac{3\bar{g}}{4\pi v_F} \; \frac{2\omega_m}{\sqrt{\gamma|\omega_m| + \left(\frac{2k_\parallel \alpha}{1+\alpha^2}\right)^2} - \left|\frac{2k_\parallel \alpha}{1+\alpha^2}\right|}, \tag{3}$$

where $\gamma = 2N\bar{g}/(\pi v_x v_y)$ and $k_\parallel$ is a deviation from a hot spot along the FS. Right at a hot spot $\Sigma(\omega_m, 0) = \omega_m \sqrt{\omega_0/|\omega_m|}$, where $\omega_0 = (9\bar{g}/(16\pi N))(2v_x v_y/v_F^2)$. In (2), (3), we neglected terms with $n \geq 2$ loops, which are small in $1/N$, and terms with $n \geq 6$ which break $1/N$ expansion [18,19]. Superconducting T_c is only weakly affected by two-loop corrections, even when $N = 1$ (Refs. [9,23]), and we expect that corrections from higher loops will remain numerically small.

Pairing vertex We add to the action the anomalous term $\Phi_0(k)\psi_{k,\alpha}(i\sigma^y)_{\alpha\beta}\psi_{-k,\beta}$ and use Eq. (1) to renormalize it into the full $\Phi(k)$. At T_c, the pairing susceptibility $\chi_{pp}(k) = \Phi(k)/\Phi_0$ must diverge for all k. The bare Φ_0 can be set constant within a patch, but has to change signs between patches separated by $\mathbf{Q}$ (the pairing symmetry at the onset of SDW order is $d-$wave [24]). The one-loop renormalization of $\Phi(k)$ at $k = (\omega \sim T, 0)$was obtained by MS:

$$\Phi(\omega \sim T, 0) = \Phi_0(1 + \lambda \log^2 \Lambda/T), \quad \lambda = \frac{\alpha}{\pi(1+\alpha^2)}, \tag{4}$$

where Λ is the smaller of ω_0 and $\alpha^4 E_F^2/\omega_0$. Notice that neither the coupling constant $\bar{g}$ not $1/N$ are present in (4), the only parameter is the ratio of the velocities α, which is a geometrical property of the FS. For a cuprate-like FS, $\alpha \sim 1$, i.e., the pairing coupling constant $\lambda = O(1)$. To understand the physics of the pairing at the QCP, it is, however, advantageous to keep λ as a free parameter and analyze the pairing at both small and moderate λ.



FIG. 1: Diagrammatic representation for the pairing vertex. The shaded triangle is the full Φ_k, the unshaded vertex is the bare Φ_0, solid lines are full fermionic propagators, and the wavy line is the Landau-overdamped spin propagator. The pairing vertex contains $i\sigma^y_{\alpha,\beta}$, the vertices where wavy and solid lins meet contain $\sigma_{\gamma\delta}$.

Let's first see where $\log^2$ renormalization comes from. The one-loop diagram for Φ contains two fermionic propagators $G(k)$ and $G(-k)$ and and one bosonic $\chi(k)$ (Fig.1). Large N allows one to restrict $\chi(\Omega_m, \mathbf{k})$ to momenta connecting points at the FS and integrate over momenta transverse to the FS in the fermionic propagators only. Because Σ does not depend on this momentum, the integration is straightforward, and yields, to logarithmic accuracy $\int GG\chi \propto \int dk_\parallel \int_T d\Omega_m (\chi(\Omega_m, k_\parallel)/|\Omega_m + \Sigma(\Omega_m, k_\parallel))|$. At $k_\parallel^2 > \gamma\Omega_m$ and $|k_\parallel| < k_F\bar{g}/v_F$, $1/|\Omega_m + \Sigma(\Omega_m, k_\parallel)|$ scales as $|k_\parallel/\Omega_m|$ and $\chi(\Omega_m, k_\parallel) \propto 1/k_\parallel^2$. Integrating over $k_\parallel$ we obtain $\int_{\gamma|\Omega_m|} dk_\parallel^2/k_\parallel^2 \propto \log|\Omega_m|$, and the remaining integral over frequency yields $\int GG\chi \propto \int_T (d\Omega_m/|\Omega_m|)\ \log|\Omega_m| \propto \log^2 T$. We see that the $\log^2 T$ dependence originates from extra logarithm from $k-$integration. This in turn is the consequence of $\Omega_m/k_\parallel$ form of self-energy $\Sigma(\Omega_m, k_\parallel)$ at $k_\parallel^2 > \gamma\Omega_m$. As $\Sigma \propto \omega$ is the property of a FL, the $\log^2 T$ renormalization comes from fermions which preserve a FL behavior at a QCP. We further see that the one-loop renormalization can be interpreted as coming from the process in which fermions are exchanging quanta of an effective local $\log\Omega$ interaction. The same process determines one-loop renormalization of Φ in CSC.

The $\log^2$ analysis can be extended beyond leading order. We assume that λ is small



FIG. 2: Numerical solution of Eq. (6) at small λ. (a) The transition temperature. When λ decreases, $2\pi\lambda\log\omega_0/T_c$ approaches 1, as in Eq. (8). (b) The eigenfunction $\Phi(y)$, where $y = k_{\|}^2/(\pi T\gamma)$. Solid and dashed lines are numerical and analytical solutions of Eq. 6, respectively. The two are very close, except for the largest $y \sim \omega_0/T$, when the cutoff becomes relevant.

but $\lambda\log^2 T = O(1)$, and sum up ladder series of $\lambda\log^2 T$ terms neglecting smaller terms at each order of loop expansion. Performing the calculations (see Supplementary material for details), we find that the the analogy with CSC does not extend beyond leading order: for CSC the summation of $\lambda\log^2 T$ terms yields $\Phi = \Phi_0/\cos[(2\lambda\log^2 T)^{1/2}]$ (Ref.[16]), and the system develops a pairing instability at $|\log T_c| = \pi/2\sqrt{2\lambda}$ (Ref. [17]). In our case, perturbation series yield $\Phi = \Phi_0 e^{\lambda\log^2 T}$, i.e., the pairing susceptibility increases with decreasing T, but never diverges.

Because the summation of the leading logarithms does not lead to a finite T_c, one has to go beyond the leading logarithmical approximation and analyze the full equation for $\Phi(k)$ at $\Phi_0 = 0$ in order to understand whether or not T_c is finite at a QCP. This is what we do next.

Full gap equation. Within our approximation, the full linearized equation for the anomalous vertex is obtained by summing up ladder diagrams and keeping the self-energy in the fermionic propagator. Integrating the r.h.s. of this equation over momenta transverse to the FS, we obtain

$$\Phi(\omega_m, k_\parallel) = \frac{3\bar{g}}{2v_F} T \sum_{m'} \int \frac{dk'_\parallel}{2\pi} \frac{\Phi(\omega_{m'}, k'_\parallel)}{|\omega_{m'} + \Sigma(\omega_{m'}, k'_\parallel)|}$$
$$\times \frac{1}{k_\parallel^2 + k_\parallel'^2 - 2e k_\parallel k'_\parallel + \gamma |\omega_m - \omega_{m'}|} \tag{5}$$

where $e = (1-\alpha^2)/(1+\alpha^2)$. The temperature at which the solution exists is T_c. Let's assume first that λ is small and search whether T_c is non-zero. At small λ, typical $k_\parallel^2$ are larger than typical $\gamma\omega_m$, and one can easily verify that the vertex $\Phi(\omega_m, k_\parallel)$ has a stronger dependence on $k_\parallel$ than on frequency. In this situation, one can approximate $\Phi(\omega_m, k_\parallel)$ by $\Phi(k_\parallel)$, explicitly sum up over frequency and reduce (5) to 1D integral equation in momentum. Introducing $\bar{T} = \pi T/\omega_0$ and $x = k_\parallel^2/(\gamma\omega_0\bar{T})$ and setting $\alpha = 1$ for simplicity, we obtain from (5)

$$\Phi(y) = \lambda \int_1 \frac{dx}{x+y} \frac{\log x}{2\sqrt{x\bar{T}} + 1} \Phi(x) \tag{6}$$

The term in denominator with $\sqrt{x\bar{T}}$ is a soft upper cutoff.

The r.h.s. of (6) contains $\log^2$ contributions from the range $x \gg y$, but, as we just found, they do not lead to a pairing instability. We therefore focus on the contribution from $x \sim y$. Because the kernel is logarithmical, we search for $\Phi(x)$ in the form $\Phi(x) = \exp[-f(p(x))]$, where $p(x) = 2\pi\lambda \log x$. Substituting this into (6), we find that the form is reproduced at $1 \ll x \ll 1/\bar{T}$, when soft cutoff can be omitted. The self-consistency condition yields (see Supplementary material)

$$f(z) = \frac{1}{2\pi^2\lambda}\left(z \arcsin z + \sqrt{1-z^2} - 1\right). \tag{7}$$

At small λ, the soft cutoff can be replaced by the boundary condition that $df(z)/dz$ must be at a maximum at $z = 2\pi\lambda|\log\bar{T}|$ (i.e., that $\Phi(y)$ rapidly drops at $x \sim 1/\bar{T}$). This condition selects $T = T_c$ to be

$$T_c \sim \omega_0 e^{-1/(2\pi\lambda)} \tag{8}$$

To verify this reasoning, we solved Eq. (6) numerically. We found very good agreement with analytical results (see Fig.2).

We see from (8) that T_c is non-zero already at infinitesimally small λ, like in BCS theory. The analogy is not accidental as the pairing predominantly comes from momenta away from hot spots, for which $x \sim \bar{T}$, i.e., $k_\| \sim (\gamma\omega_0)^{1/2} \sim k_F \bar{g}/E_F$. For $\bar{g} \ll E_F$, when the calculations are under control, maximal $k_\|$ is still much smaller than k_F, i.e., the pairing is confined to a region near a hot spot. At the same time, typical $\gamma\omega \geq \gamma T_c$ are much smaller then $\gamma\omega_0$, fermionic self-energy for $k_\| \sim (\gamma\omega_0)^{1/2}$ has the FL form ($\Sigma(\omega_m, k_\|) \propto \omega_m/|k_\||$), and for $x \sim y$ in (6) the momentum integration does not give rise to an additional logarithm. The instability at T_c is then a conventional Cooper instability of a FL with a weak and non-singular attractive interaction. In other words, for small λ, the pairing at a SDW QCP is entirely a FL phenomenon.

A non-zero T_c at small λ is the consequence of the dependence of the self-energy on the momenta along the FS. Earlier works [8,9] neglected this momentum dependence and approximated the self-energy by its non-FL form $\Sigma(\omega) = \omega_m(\omega_0/|\omega_m|)^{1/2}$ at a hot spot. These studies found a different result: T_c becomes non-zero only if the coupling exceeds a certain threshold, like in the pairing problem at a QCP at $q = 0$ (Refs.[14,25]. Specifically, because $\Sigma = \Sigma(\omega_m)$, the anomalous vertex Φ also depends only on frequency, and Eq. (5) reduces to 1D integral equation in frequency rather than in momentum:

$$\Phi(\omega_m) = \pi^2 \lambda T \sum_{m'} \frac{\Phi(\omega_{m'})}{\sqrt{|\omega_{m'}|} Z_{\omega_{m'}} \sqrt{|\omega_m - \omega_{m'}|}}. \tag{9}$$

where $Z_{\omega_{m'}} = 1 + \sqrt{|\omega_{m'}|/\omega_0}$. This equation has been solved for arbitrary λ [8], and the result is that T_c becomes non-zero only when λ exceeds a critical value $\lambda_c = 0.035$. Near critical coupling $T_c \sim \omega_0 e^{-1.36/(\lambda-\lambda_c)^{1/2}}$), and for physical $\lambda = 1/2\pi$, $T_c = 0.17\omega_0$. Note that the pairing in this case comes from fermions with non-FL self-energy, yet the overall scale for T_c is the same ω_0 as in Eq. (8).

T_c **at moderate coupling.** For a generic 2D FS, v_x and v_y are comparable, and λ from (4) is of order one. We solved Eq. 5 for T_c numerically for $\alpha = 1$, when $\lambda = 1/(2\pi)$, and found

$$T_c \approx 0.04\omega_0. \tag{10}$$

Typical $k_\|^2 \sim \gamma\omega_0$ and typical $\gamma\omega \sim \gamma T_c$ are now comparable, i.e., the pairing comes from fermions for which self-energy is in a grey area between a FL and a non-FL. Numerically T_c is still small compared to ω_0, i.e., typical $k_\|^2$ are still larger than typical $\gamma\omega$. To a good accuracy the self-energy of typical fermions can still be approximated by the FL form, i.e., even at $\lambda = O(1)$ the pairing at a SDW QCP is still predominantly a FL phenomenon.



FIG. 3: Numerical solution of Eq. (5) for various $\alpha = v_y/v_x$ (the Fermi velocities at hot spots separated by (π,π) are $\mathbf{v}_{F,1} = (v_x, v_y)$ and $\mathbf{v}_{F,2} = (-v_x, v_y)$. We set $\bar{g}a/v_F \approx 0.032$. For $\alpha = O(1)$ the momentum dependence of the fermionic self-energy dominates, pairing is a FL phenomenon, and $T_c \approx 0.04\omega_0$. At smaller α, when $\bar{g} > \alpha^{3/2} v_F/a$ (to the left of vertical dashed line), frequency dependence of the self-energy prevails, the pairing comes from fermions with non-FL self-energy, and $T_c \approx 0.17\omega_0$, up to corrections of order $(\bar{g}a/v_F)^{1/2}$. In our case, these corrections reduce this number from 0.17 toto 0.15.

Eq. (10) agrees with the numerical solution of the gap equation along the full FS, without restriction to hot spots [22]. In notations of Ref. [22], $T_c = v_F/af(u)$, where dimensionless $u = 4\omega_0 a/(3v_F)$. Eq. (10) implies that $f(u) = 0.03u$ at small u. This agrees well with the numerical solution. According to various numerics [26], at larger $u \geq 1/2$, $f(u)$ saturates at a value around $0.015 - 0.02$ and at even larger u decreases as $1/u$ because of Mott physics.

At strong nesting, when $\alpha \ll 1$ and $\gamma \propto 1/\alpha$ is large, the pairing eventually become determined by fermions whose self-energy has a non-FL form. This happens when $(\gamma\omega_0)^{1/2}$ becomes larger than maximal possible $|k_\parallel|$ along a Fermi arc, which is of order $1/a$, i.e., when $v_F/a \gg \bar{g} > \alpha^{3/2} v_F/a$. In this situation, the momentum dependence of $\Sigma(\omega_m, k_\parallel)$ becomes irrelevant at $\omega_m \sim \omega_0$, $\Phi(\omega_m, k_\parallel)$ becomes predominantly frequency-dependent, and T_c recovers the value $0.17\omega_0$, which is T_c for momentum-independent self-energy. We verified numerically that this is indeed the case. We plot T_c vs α in Fig. 3. At $\alpha = O(1)$, $T_c \approx 0.04\omega_0$, and at small enough α T_c approaches $0.017\omega_0$. We caution, however, that the limit $\alpha \ll 1$ has to be taken with care as nesting may generate additional singularities at higher-loop orders.

Conclusions. In this paper we analyzed the equation for superconducting T_c at the onset of SDW order. We demonstrated that the leading perturbation correction to the bare pairing vertex contains $\log^2 T$, but the series of $\log^2 T$ renormalizations do not give rise to the pairing instability. Yet, T_c is finite, even when coupling is small, because of subleading, $\log T$ terms. We showed that the pairing predominantly comes from fermions which retain a FL form of the self-energy at a QCP, i.e., the pairing at a SDW QCP is actually a FL phenomenon. The overall scale of T_c is set by the interaction ($\omega_0 \sim \bar{g}$), as long as the interaction is smaller than the Fermi energy.

The large N approximation which we used to justify the gap equation requires caution because (i) two-loop $1/N$ corrections are logarithmically singular [9,23], and (ii) diagrams with $n \geq 6$ loops are not small in $1/N$ and may give rise to additional logarithms [18,19,21]. On the other hand, the fact that our T_c is of order ω_0, which is the upper limit of logarithmical behavior, likely implies that logarithmical renormalizations, not included into our analysis, do not change T_c qualitatively and do not affect the conclusion that the pairing at SDW QCP is a FL phenomenon.

We acknowledge stimulating discussions with Y.B. Kim, S.S. Lee, M.A. Metlitski, S. Sachdev, T. Senthil, and A-M Tremblay. The research has been supported by DOE DE-FG02-ER46900.

---

[1] S. Sachdev and B. Keimer, Physics Today, 64, 29, (2011).

[2] K. Hashimoto et al., Science 336, 1554 (2012).

[3] A. Chubukov, D. Pines and J. Schmalian *A Spin Fluctuation Model for D-wave Superconductivity* in 'The Physics of Conventional and Unconventional Superconductors' edited by K.H. Bennemann and J.B. Ketterson (Springer-Verlag), 2002.

[4] H. v. Löhneysen, A. Rosch, M. Vojta, and P. Wölfle, Rev. Mod. Phys. **79**, 1015 (2007).

[5] E. Fradkin, S. A. Kivelson, M. J. Lawler, J. P. Eisenstein, A. P. Mackenzie, Annual Reviews of Condensed Matter Physics **1**, 153 (2010).

[6] P. A. Lee, Phys. Rev. Lett. **63**, 680 (1989); J. Polchinski, Nucl. Phys. B **422**, 617 (1994); Y.-B. Kim, A. Furusaki, X.-G. Wen, and P. A. Lee, Phys. Rev. B **50**, 17917 (1994); C. Nayak and F. Wilczek, Nucl. Phys. B 417, 359 (1994); 430, 534 (1994); B. L. Altshuler, L. B. Ioffe, and A. J. Millis, Phys. Rev. B **50**, 14048 (1994); **52**, 5563 (1995); S. Chakravarty, R. E. Norton, and O. F. Syjuasen, Phys. Rev. Lett. 74, 1423 (1995); C.J. Halboth and W. Metzner, Phys. Rev. Lett. **85**, 5162 (2000); J. Quintanilla and A. J. Schofield, Phys. Rev. B **7**4, 115126 (2006); J. Rech, C. Pépin, and A. V. Chubukov, Phys. Rev. B **74**, 195126 (2006); T. Senthil, Phys. Rev. B **78**, 035103 (2008); M. Zacharias, P. Wölfle, and M. Garst, Phys. Rev. B **80**, 165116 (2009); J. She and J. Zaanen, Phys. Rev. B 80, 184518 (2009); D.L. Maslov and A.V. Chubukov, Phys. Rev. B **81**, 045110 (2010). M.A. Metlitski and S. Sachdev, Phys. Rev. B 82, 075127 (2010); D. F. Mross, J. McGreevy, H. Liu, and T. Senthil, Phys. Rev. B 82, 045121 (2010); Casper Drukier, Lorenz Bartosch, Aldo Isidori, and Peter Kopietz Phys. Rev. B 85, 245120 (2012).

[7] A. J. Millis, S. Sachdev, and C. M. Varma, Phys. Rev. B 37, 4975 (1988).

[8] Ar. Abanov, A. V. Chubukov, and A.M. Finkelstein, Europhys. Lett. **54**, 488 (2001).

[9] A. Abanov, A. V. Chubukov, and J. Schmalian, Adv. Phys., **52**, 119 (2003).

[10] E.G. Moon and A.V. Chubukov, J. Low Temp. Phys., 161, 263-281 (2010).

[11] E. G. Moon, and S. Sachdev, Phys. Rev. B **80**, 035117 (2009).

[12] J.-H. She, B. J. Overbosch, Y.-W. Sun, Y. Liu, K. E. Schalm, J. A. Mydosh, and J. Zaanen, Phys. Rev. B 84, 144527 (2011).

[13] N. E. Bonesteel, I. A. McDonald, and C. Nayak, Phys. Rev. Lett. 77, 3009 (1996).

[14] Ar. Abanov, B. Altshuler, A. Chubukov, and E. Yuzbashyan, unpublished.

[15] D. V. Khveshchenko and W. F. Shively, Phys. Rev. B 73, 115104 (2006); A. V. Chubukov and A. M. Tsvelik, Phys. Rev. B 76, 100509 (2007).

[16] A. Chubukov, and J. Schmalian, Phys. Rev. B. **72**, 174520 (2005).

[17] D. T. Son, Phys. Rev. D. **59**, 094019 (1999).

[18] S.-S. Lee, Phys. Rev. B **80**, 165102 (2009).

[19] M.A. Metlitski and S. Sachdev, Phys. Rev. B 82, 075128 (2010).

[20] A.J. Millis, Phys. Rev. B 45, 13047 (1992).

[21] S. A. Hartnoll, D. M. Hofman, M. A. Metlitski, and S. Sachdev Physical Review B 84, 125115 (2011); E. Berg, M. A. Metlitski, and S. Sachdev, arXiv:1206.0742.

[22] Ar. Abanov, A.V. Chubukov, and M. Norman, Phys. Rev. B 78, 220507 (2008).

[23] Ar. Abanov and A.V. Chubukov, Phys. Rev. Lett., **93**, 255702 (2004).

[24] D. Scalapino, Phys. Rep., **250**, 329 (1995).

[25] M.A Metlitski et al, unpublished.

[26] P. Monthoux and D. J. Scalapino, Phys. Rev. Lett. 72, 1874 (1994); T. Dahm and L. Tewordt, Phys. Rev. B 52, 1297 (1995); D. Manske, I. Eremin and K. H. Bennemann, Phys. Rev. B 67, 134520 (2003); St. Lenck, J. P. Carbotte and R. C. Dynes, Phys. Rev. B 50, 10149 (1994); B. Kyung, J.-S. Landy, A.-M. S. Tremblay, Phys. Rev. B 68, 174502 (2003); T. Maier et al., Phys. Rev. Lett. 95, 237001 (2005); K. Haule and G. Kotliar, Phys. Rev. B 76, 104509 (2007); S. S. Kancharla et al., Phys. Rev. B 77, 184516 (2008); D. Senechal and A-M. S. Tremblay, Phys. Rev. Lett. 92, 126401 (2004).

I. SUPPLEMENTARY MATERIAL

A. Perturbation Theory

We add a fictitious anomalous term $\Phi_0 \psi_{k,\alpha}(i\sigma^y)_{\alpha\beta}\psi_{-k,\beta}$ to the original Lagrangian to generate a bare pairing vertex, renormalize it by particle-particle interaction, and obtain the pairing susceptibility as a ratio of the fully renormalized and bare pairing vertices. Within the approximations which we discuss in the main text, the diagrams for the renormalization of the pairing vertex form ladder series, shown in Fig. 4, however each line is the full fermionic Green's function, which includes the self-energy.



FIG. 4: Perturbation expansion for the pairing vertex.

The renormalized vertex Φ depends on fermionic frequency ω_m, fermionic momentum along the Fermi surface, $k_\|$, and on temperature. At high enough temperatures $\Phi(\omega_m, k_\|, T)$ weakly deviates from Φ_0, but at T_c the ratio Φ/Φ_0 should diverge for all frequencies and momenta. For definiteness, we set $k_\|$ to zero and ω_m to πT, i.e., consider $\Phi(\pi T, 0, T) = \Phi(T)$. To simplify the formulas, we also set the ratio of Fermi velocities $\alpha = v_y/v_x$ to one.

Consider one-loop renormalization of Φ (the diagram with one wavy line in Fig. 4). After integration over momenta transverse to the Fermi surface (FS) we obtain, replacing the summation over ω_m by integration over $|\omega_m| > \pi T$,

$$\Phi(T) = \Phi_0 \left(1 + \pi(\omega_0\gamma)^{1/2} \int \frac{d\omega_m dk_\|}{4\pi^2} \frac{1}{|\omega_m + \Sigma(\omega_m, k_\|)|} \frac{1}{k_\|^2 + \gamma|\omega_m|}\right) \tag{11}$$

where, we remind, $\Sigma(\omega_m, k_\|) = \text{sign}\ \omega_m |\Sigma(\omega_m, k_\|)|$, and [9,19]

$$|\Sigma(\omega_m, k_\|)| = \sqrt{\omega_0}\left(\sqrt{|\omega| + k_\|^2/\gamma} - |k_\||/\sqrt{\gamma}\right). \tag{12}$$

Rescaling $k_\| = (|\omega|\gamma)^{1/2} z$ we obtain

$$\Phi(T) = \Phi_0 \left(1 + \lambda \int_{\pi T}^{\infty} \frac{d\omega_m}{\omega_m} \int dz \frac{\sqrt{z^2+1} + |z|}{z^2+1} \frac{1}{1 + \left(\frac{\omega_m}{\omega_0}\right)^{1/2} (\sqrt{1+z^2} + |z|)}\right) \tag{13}$$

where $\lambda = 1/(2\pi)$. One can easily make sure that the the integral in the r.h.s. of (13) contains $\log^2 \omega_0/T$, which comes from large z, i.e., from $k_\parallel^2 \gg |\omega_m|\gamma$ (Ref.[19]), and $\log \omega_0/T$, which comes from $z = O(1)$. The coupling constant λ is a fixed number, but in the spirit of perturbation theory we assume that λ is a small parameter. Then, obviously, $\lambda \log^2 \omega_0/T$ is more relevant than $\lambda \log \omega_0/T$, i.e., to logarithmic accuracy [19]

$$\Phi(T)\Phi_0 \left(1 + \lambda \log^2 \omega_0/T\right) \tag{14}$$

We now check whether the series of $\lambda \log^2 \omega_0/T$ give rise to a pairing instability at $\log \omega_0/T \sim 1/\sqrt{\lambda}$. Because $\log^2 \omega_0/T$ comes from $k_\parallel^2 \gg |\omega_m|\gamma$, we can expand the self-energy in $\gamma|\omega|/k_\parallel^2$, i.e., approximate $|\Sigma(\omega_m, k_\parallel)|$ in (12) by

$$|\Sigma(\omega_m, k_\parallel)| \approx \frac{\sqrt{\omega_0\gamma}}{2} \frac{\omega_m}{|k_\parallel|} \tag{15}$$

Using this self-energy, we obtain at two-loop order ($k_{\parallel,1} = k_1, k_{\parallel,2} = k_2$)

$$\Phi(T) = \Phi_0 \left(1 + \lambda \log^2 \omega_0/T + \lambda^2 \int_{\sqrt{\gamma\pi T}}^{\sqrt{\gamma\Lambda}} \frac{2k_1 dk_1}{k_1^2} \int_{\pi T}^{k_1^2/\gamma} \frac{d\omega_1}{\omega_1} \int_{\sqrt{\gamma\pi T}}^{\sqrt{\gamma\Lambda}} \frac{2k_2 dk_2}{k_2^2 + k_1^2} \int_{\pi T}^{k_2^2/\gamma} \frac{d\omega_2}{\omega_2} + O(\lambda^3)\right) \tag{16}$$

Evaluating the integrals we find

$$\Phi(T) = \Phi_0 \left(1 + \lambda \log^2 \omega_0/T + \frac{1}{2}\lambda^2 \log^4 \omega_0/T + ...\right) \tag{17}$$

To understand what are the prefactors from higher-order terms, we note that $\log^4$ term comes from the region of internal momenta $k_2 \gg k_1$, or, more specifically, the momentum k_1 in the cross-section which is farther from the vertex sets the lower cutoff of momentum integration over k_2 in the cross-section, which is closer to the vertex. This sheds light on the general pattern – we verified that at order M the largest, $\log^{2M} \omega_0/T$ term comes from comes $k_M > k_{M-1} > ... > k_1$, where k_M is the momentum in the cross-section, which is the closest to the vertex. The same trend was earlier detected in the perturbation theory for the pairing vertex in 2D systems for which fermionic self-energy depends only on frequency [8]. A simple exercise in combinatorics then yields, for the full series

$$\begin{aligned}\Phi(T) &= \left(1 + \lambda \log^2 \omega_0/T + \frac{1}{2}\lambda^2 \log^4 \omega_0/T + \frac{1}{6}\lambda^3 \log^6 \omega_0/T + ...\right) \\ &= \Phi_0 \sum_{M=0}^{\infty} \frac{\lambda^M}{M!} \log^{2M} \frac{\omega_0}{T} = \Phi_0 e^{\lambda \log^2 \omega_0/T}\end{aligned} \tag{18}$$

We see that the calculation to $\log^2$ accuracy does not give rise to a pairing instability: the pairing susceptibility, Φ/Φ_0 increases with decreasing T, but does not diverge at any finite T.

1. *A comparison with color superconductivity*

The presence of $\log^2 \omega_0/T$ in perturbation theory brings in the comparison with the problem of color superconductivity (CSC), where perturbative expansion also contains series of $\lambda \log^2 T$ terms. For that problem, however, previous works have demonstrated that the summation of $\log^2$ terms does lead to a pairing instability. It is therefore instructive to compare perturbation theory for CSC problem with our case and see where the two cases differ.

In CSC problem [16,17], as well as in condensed-matter problems of the pairing mediated by gapless collective excitations in three spatial dimensions [10], fermionic self-energy depends only on frequency (and is actually irrelevant to the pairing problem), and the extra logarithm, in addition to a Cooper one, appears because the effective interaction, integrated over momenta along the FS, has logarithmic dependence on frequency. Because the integration over momenta $k_{\|}$ can be done independently in any cross-section in Fig. 4, the pairing can be analyzed within the effective local model of fermions with dynamical interaction [17]

$$\chi(\omega_m) = \lambda \log \frac{\Lambda}{|\omega_m|} \tag{19}$$

Substituting this form of the interaction (wavy line) into the one-loop diagram for Φ, we obtain

$$\Phi(T) = \Phi_0 \left(1 + 2\lambda \int_{\pi T}^{\Lambda} \frac{d\omega}{\omega} \log \frac{\Lambda}{|\omega|}\right) = \Phi_0 \left(1 + \lambda \log^2 \frac{\Lambda}{\pi T}\right) \tag{20}$$

This is the same result as in our case. At two-loop order, however, the prefactor for $\lambda^2 \log^4 \Lambda/T$ term is different from the one in our case. For CSC we have, at two-loop order

$$\Phi(T) = \Phi_0 \left(1 + \lambda \log^2 \frac{\Lambda}{\pi T} + \lambda^2 \int_{\pi T}^{\Lambda} \frac{2d\omega'}{\omega'} \log \frac{\Lambda}{|\omega'|} \int_{\pi T}^{\Lambda} \frac{2d\omega''}{\omega''} \log \frac{\Lambda}{|\omega' - \omega''|}\right) \tag{21}$$

It is natural to divide the integration range over ω'' into two regimes, $\omega'' \gg \omega'$ and $\omega'' \ll \omega'$.

It turns out that *both* regimes give contributions of order $\log^4 \Lambda/T$. We have

$$
\begin{aligned}
&\lambda^2 \int_{\pi T}^{\Lambda} \frac{2d\omega'}{\omega'} \log\frac{\Lambda}{|\omega'|} \int_{\pi T}^{\Lambda} \frac{2d\omega''}{\omega''} \log\frac{\Lambda}{|\omega' - \omega''|} \\
=&\lambda^2 \int_{\pi T}^{\Lambda} \frac{2d\omega'}{\omega'} \log\frac{\Lambda}{|\omega'|} \left(\int_{\pi T}^{\omega'} \frac{2d\omega''}{\omega''} \log\frac{\Lambda}{\omega'} + \int_{\omega'}^{\Lambda} \frac{2d\omega''}{\omega''} \log\frac{\Lambda}{\omega''} \right) \\
=&\lambda^2 \left(2\int_{\pi T}^{\Lambda} \frac{2d\omega'}{\omega'} \log\frac{\Lambda}{|\omega'|} \log\frac{\omega'}{\pi T} \log\frac{\Lambda}{\omega'} + \int_{\pi T}^{\Lambda} \frac{2d\omega'}{\omega'} \log\frac{\Lambda}{|\omega'|} \log^2\frac{\Lambda}{\omega'} \right) \\
=&\lambda^2 \left(\frac{1}{3} \log^4\frac{\Lambda}{\pi T} + \frac{1}{2} \log^4\frac{\Lambda}{\pi T} \right) \\
=&\frac{5}{6}\lambda^2 \log^4\frac{\Lambda}{\pi T}, \qquad (22)
\end{aligned}
$$

Combining one-loop and two-loop terms, we obtain

$$
\Phi(T) = \Phi_0 \left(1 + \lambda \log^2\frac{\Lambda}{\pi T} + \frac{5}{6}\lambda^2 \log^4\frac{\Lambda}{\pi T} \right) \qquad (23)
$$

which is different from the two-loop result in our case, Eq. (17). The difference comes about because for CSC there is no requirement that highest power of the logarithm comes from particular hierarchy of running frequencies – at two loop order, there is $\log^4$ contribution from the range where the highest frequency in the cross-section next to the vertex, and from the range when the highest frequency is in the cross-section farthest from the vertex.

The series of $\lambda \log^2$ terms for CSC problem have been summed up in Ref. [16], an the result is

$$
\begin{aligned}
\Phi(\omega = \pi T) &= \Phi_0 \left(1 + \lambda \log^2\frac{\Lambda}{\pi T} + \frac{5}{6}\lambda^2 \log^4\frac{\Lambda}{\pi T} + O(\lambda^3) \right) \\
&= \frac{\Phi_0}{\cos[(2\lambda \log^2 \Lambda/\pi T)^{1/2}]} \qquad (24)
\end{aligned}
$$

B. Linearized gap equation

1. *Interplay between characteristic momenta and frequency*

We start with the Eq (5) in the main text. For $\alpha = 1$ we have

$$
\Phi(\omega_m, k_\parallel) = \pi \left(\omega_0 \gamma\right)^{1/2} T \sum_{m'} \int \frac{dk'_\parallel}{2\pi} \frac{\Phi(\omega_{m'}, k'_\parallel)}{|\omega_{m'} + \Sigma(\omega_{m'}, k'_\parallel)|} \times \frac{1}{k_\parallel^2 + k'^2_\parallel + \gamma|\omega_m - \omega_{m'}|} \qquad (25)
$$

where the self-energy Σ is given by (12). The dependence on system parameters ω_0 and γ can be eliminated if we measure ω and T in units of ω_0, and measure $k_\parallel$ in units of $(\omega_0\gamma)^{1/2}$.

Introducing $\bar{\omega} = \omega/\omega_0$, $\bar{T} = T/\omega_0$, and $\bar{k}_\parallel = k_\parallel/(\omega_0\gamma)^{1/2}$, we re-write (25) as

$$\Phi(\bar{\omega}_m, \bar{k}_\parallel) = 2\pi^2\lambda\bar{T}\sum_{m'}\int\frac{d\bar{k}'_\parallel}{2\pi}\frac{\Phi(\bar{\omega}_{m'}, \bar{k}'_\parallel)}{|\bar{\omega}_{m'} + \bar{\Sigma}(\bar{\omega}_{m'}, \bar{k}'_\parallel)|} \times \frac{1}{\bar{k}^2_\parallel + \bar{k}'^2_\parallel + |\bar{\omega}_m - \bar{\omega}_{m'}|}, \tag{26}$$

where, as before, $\lambda = 1/(2\pi)$, and

$$\bar{\Sigma}(\bar{\omega}_m, \bar{k}_\parallel) = \left(\sqrt{|\bar{\omega}| + \bar{k}^2_\parallel} - |\bar{k}_\parallel|\right) \tag{27}$$

In this section we again consider λ not as a given constant but as a free parameter, and analyze the limit $\lambda \ll 1$. The question we address is whether T_c is non-zero already at arbitrary small λ, or it only emerges when λ exceeds a certain threshold.

Because λ is treated as small parameter, $\bar{T}_c$ is expected to be small, and to get the pairing we need to explore logarithmical behavior which comes from frequency scale larger than T_c. Accordingly, we replace $\bar{T}\sum_{m'}$ by $(1/2\pi)\int d\omega_{m'}$ and set $\pm\pi T$ as the lower limits of the integration over positive and negative $\omega_{m'}$, respectively. We then have, instead of (26)

$$\Phi(\bar{\omega}_m, \bar{k}_\parallel) = \pi\lambda\int d\bar{\omega}_{m'}\int\frac{d\bar{k}'_\parallel}{2\pi}\frac{\Phi(\bar{\omega}_{m'}, \bar{k}'_\parallel)}{|\bar{\omega}_{m'} + \bar{\Sigma}(\bar{\omega}_{m'}, \bar{k}'_\parallel)|} \times \frac{1}{\bar{k}^2_\parallel + \bar{k}'^2_\parallel + |\bar{\omega}_m - \bar{\omega}_{m'}|}, \tag{28}$$

In general, $\Phi(\bar{\omega}_m, \bar{k}_\parallel)$ is a function of both arguments, but in proper limits the dependence on one of the arguments is stronger than on the other. In the main text we consider the limits $\bar{k}^2_\parallel \gg \bar{\omega}_m$ and $\bar{k}^2_\parallel \ll \bar{\omega}_m$. In the first case, the momentum dependence is stronger than frequency dependence, and $\Phi(\bar{\omega}_m, \bar{k}_\parallel)$ can be approximated by $\Phi(\bar{k}_\parallel)$. The fermionic self-energy at $\bar{k}^2_\parallel \gg \bar{\omega}_m$ behaves as

$$\bar{\Sigma}(\bar{\omega}_m, \bar{k}_\parallel) \approx \frac{|\bar{\omega}_m|}{2|\bar{k}_\parallel|} \tag{29}$$

Substituting this form into (28) we obtain

$$\Phi(\bar{k}_\parallel) = 4\lambda\int\frac{\bar{k}'_\parallel d\bar{k}'_\parallel}{\bar{k}^2_\parallel + \bar{k}'^2_\parallel}\frac{1}{2\bar{k}'_\parallel + 1}\Phi(k'_\parallel)\int_{\bar{T}}^{\bar{k}^2_\parallel}\frac{d\bar{\omega}'_m}{\bar{\omega}'_m}. \tag{30}$$

Integrating explicitly over ω_m and introducing new variable $x = \bar{k}'^2_\parallel/\bar{T}$, we obtain Eq 6 in the main text.

$$\Phi(y) = 2\lambda\int_1\frac{dx}{x+y}\frac{\log x}{2\sqrt{x\bar{T}} + 1}\Phi(x), \tag{31}$$

In the opposite limit, $\bar{k}^2_\parallel \ll \bar{\omega}$, the momentum dependence of the self-energy is small, and $\bar{\Sigma}$ can be approximated by its value at the hot spot, i.e., $\bar{\Sigma} \approx \sqrt{|\bar{\omega}_m|}$. This limit can only be justified at small or large ratio of v_y/v_x because otherwise substituting this $\bar{\Sigma}$ into (28) we obtain that typical internal $\bar{k}_{\|}'^2$ are of order $\bar{\omega}'$. Nevertheless, if we assume that the momentum dependence of the fermionic self-energy can be neglected, at least for order-of-magnitude estimates, we obtain that $\Phi(\bar{\omega}_m, \bar{k}_{\|})$ can be approximated by $\Phi(\bar{\omega}_m)$, and the gap equation becomes [8,9]

$$\Phi(\bar{\omega}_m) = \pi^2 \lambda \bar{T} \sum_{m'} \frac{\Phi(\bar{\omega}_{m'})}{\sqrt{|\bar{\omega}_{m'}|}\sqrt{|\bar{\omega}_m - \bar{\omega}_{m'}|}(1 + \sqrt{|\bar{\omega}_{m'}|})}. \tag{32}$$

This is Eq. (9) in the main text.

2. *Solution of the Eq. 6*

We first replace the soft cutoff imposed by $\left(2\sqrt{x\bar{T}} + 1\right)^{-1}$ with a hard cutoff,

$$\Phi(y) = 2\lambda \int_1^{1/\bar{T}} \frac{dx}{x+y} \log x \; \Phi(x) \tag{33}$$

The integration over $x \gg y$ gives rise to $\log^2$ terms in the perturbation theory, which, as we know from the analysis in the previous section, do not give rise to the pairing instability. We therefore focus on the range $x \sim y$. The contribution from this range gives rise to a single logarithm ($\log 1/\bar{T}$) if we momentary assume that $\Phi(x)$ is a constant. Like we did before, we treat λ as a small parameter and check whether Eq. (33) has a solution at a finite $\bar{T}$.

For small λ, $\bar{T}$ is expected to be also small, i.e., the upper limit of the integration in (33) is a large number. We first consider y and $x \sim y$ in the range $1 \ll y, x \ll 1/\bar{T}$, and search for the solution of (33) in the form $\Phi(x) = \exp[-f(p(x))]$, where $p(x) = 2\pi\lambda \log x$. Plugging this function back into the equation and introducing a new variable $z = x/y$, we obtain,

$$e^{-f(2\pi\lambda \log y)} = 2\lambda \log y \int_{\frac{1}{y}}^{\infty} \frac{dz}{z+1} \left(1 + \frac{\log z}{\log y}\right) e^{-f(2\pi\lambda(\log y + \log z))} \tag{34}$$

We assume and then verify that typical z are of order one, i.e., $\log z \ll \log y$. Taylor expanding in $\log z / \log y$ we obtain

$$\begin{aligned} 1 =& 2\lambda \log y \int_0^{\infty} \frac{dz}{z+1} \frac{1}{z^{Q(2\pi\lambda \log y)}} \\ =& 2\lambda \log y \; \frac{\pi}{\sin(\pi Q(2\pi\lambda \log y))}. \end{aligned} \tag{35}$$

where we introduced

$$Q(p) \equiv \frac{df(p)}{dp}. \tag{36}$$

Solving (35) we obtain

$$Q(2\pi\lambda \log y) = \frac{1}{\pi} \arcsin(2\pi\lambda \log y). \tag{37}$$

Integrating over $\log y$, we have

$$f(p) = \frac{1}{2\pi^2\lambda} \left[p \arcsin(p) + \sqrt{1-p^2} - 1 \right], \tag{38}$$

which is Eq 7 of the main text.



FIG. 5: Numerical solution of $Q(p)$ and its comparison with theory. In this figure we have set $\lambda = 3.7 \times 10^{-4}$.

Lastly we consider the boundary condition at $y = y_0 = 1/\bar{T}$. For such y, the upper limit is 1. Substituting formally the trial solution $\Phi(x) = \exp[-f(p(x))]$ into the r.h.s. of (33) we obtain

$$1 = 2\lambda \log y_0 \int_0^1 \frac{dz}{z+1} \frac{1}{z^{Q(2\pi\lambda \log y_0)}} \tag{39}$$

One can easily make sure that to satisfy this equation, $Q(2\pi\lambda \log y_0)$ must be *larger* than if we take our result, Eq. (37), and set $y = y_0$ there. At vanishingly small $\bar{T}$, the presence of the upper limit of the integration over x in (33) becomes relevant only for y infinitesimally close to y_0 (for smaller y, the upper limit can be safely set to infinity). This implies that at $\bar{T} \to 0$, $Q(p)$ undergoes a finite jump at $y = y_0$, hence at the actual $T = T_c$

$$\left.\frac{dQ(p)}{dp}\right|_{p=2\pi\lambda\log 1/\bar{T}_c} = \infty. \tag{40}$$

This last equation is satisfied if $2\pi\lambda \log 1/\bar{T}_c = 1$ Restoring the parameters, we then obtain Eq 8 in the main text. For this T_c, the jump in Q at the boundary is between $Q = 0.5$ (Eq. (37) at $y = y_0 = 1/\bar{T}_c$) and $Q = 0.73$, which is the solution of (39) at T_c.

In Fig. 5 we plot $Q(p)$ obtained from the numerical solution of Eq. (33) and compare it with the (approximate) analytical solution presented in this section. We see that the agreement is quite good.